

PRODUCT INFORMATION SHEET

MULTIMIN[®] EVOLUTION INJECTION FOR CATTLE

- ✔ Promotes cattle fertility and immunity
- ✔ Treats 100 head of cattle [500 kg]
- ✔ Contains 90 mg/mL of trace minerals
- ✔ Trace mineral top up for high demand periods [calving, joining, weaning]

ACTIVE CONSTITUENTS


- ✔ 60 g/L ZINC [as disodium zinc EDTA]
- ✔ 15 g/L COPPER [as disodium copper EDTA]
- ✔ 10 g/L MANGANESE [as disodium manganese EDTA]
- ✔ 5 g/L SELENIUM [as sodium selenite]

PRESENTATION

500 mL pillow pack

WHEN TO USE MULTIMIN[®] EVOLUTION

Subcutaneous injection in middle of side of neck.



ANIMAL AGE: UP TO 1 YEAR

[1 mL per 50 kg]

Liveweight (kg)	Injection Dose Rate (mL)
25	0.5
50	1.0
75	1.5
100	2.0
125	2.5
150	3.0
175	3.5
200	4.0
225	4.5
250	5.0

ANIMAL AGE: FROM 1 TO 2 YEARS

[1 mL per 75 kg
round down to the nearest 0.5 mL]

Liveweight (kg)	Injection Dose Rate (mL)
190	2.5
225	3.0
265	3.5
300	4.0
340	4.5
375	5.0
415	5.5

ANIMAL AGE: OVER 2 YEARS

[1 mL per 100 kg]

Liveweight (kg)	Injection Dose Rate (mL)
400	4.0
500	5.0
600	6.0
700	7.0
800	8.0
900	9.0
1000	10.0

BEEF CATTLE

Cows and Heifers	Weaners	Bulls	Feedlotting/backgrounding
30 Days pre-joining*	At or before weaning	90 Days pre-joining^	Before or at induction
30 Days pre-calving		At joining	

DAIRY CATTLE

Cows	Heifers	Calves
30 Days pre-joining*	30 Days pre-joining*	At birth
At or before dry off	30 Days pre-calving	At weaning

For northern cattle producers, Multimin[®] Evolution can also be used once every three months, or at each muster.

*Use Multimin[®] Evolution 6-8 weeks before artificial breeding [TAI, ET, MOET, etc].

^The beef bull program can also be applied to dairy bulls.

DOSAGE AND ADMINISTRATION:

Use the contents within 3 months of first broaching of the container, with a maximum of 3 broachings over this period. Discard the unused portion.

Use in cattle only by subcutaneous injection. Multimin[®] Evolution can be concurrently administered with other animal health products. Always administer in a different injection site to other injectable products. Do not use concurrently with vaccines, drenches, or boluses containing selenium or copper.

WITHHOLDING PERIODS:

MEAT: Zero (0) days.

MILK: Zero (0) days.

TRADE ADVICE:

EXPORT SLAUGHTER INTERVAL (ESI): Zero (0) days.